

DEGENERATE BESS AT FUTURE e^+e^- COLLIDERS *

Daniele Dominici
Dipartimento di Fisica dell'Università and
Istituto Nazionale di Fisica Nucleare
I-50125 Firenze, Italy

ABSTRACT

The sensitivity of future e^+e^- colliders to the new physics deriving from a model of strong electroweak symmetry breaking is discussed. The model considered is an effective lagrangian description of Goldstone bosons and of new vector resonances degenerate in mass.

1. Introduction

The model proposed in ¹⁾ is an effective lagrangian description of Goldstone bosons and new vector and axial vector resonances as the possible manifestation at low energy of the strong interacting sector.

To build the effective low energy theory describing Goldstones and vectors, one can use the non linear representations of a chiral symmetry G and considering (à la Weinberg) the ρ as the gauge field of the unbroken symmetry group H . In a completely equivalent way one can use the *hidden gauge symmetry* approach. Theories with non linearly realized symmetry $G \rightarrow H$ can be linearly realized by enlarging the gauge symmetry G to $G \otimes H' \rightarrow H_D = \text{diag}(H \otimes H')$. H' is a local gauge group and the ρ is the gauge field associated to H' .

* Talk presented at the IInd Rencontres du Vietnam, At the Frontiers of the Standard Model, Hochiminh Ville, October 21-28 1995. This work was partially supported by a grant under the European Human Mobility Program on "Tests of Electroweak Symmetry Breaking at Future European Colliders".

The BESS (Breaking Electroweak Symmetry Strongly) model ²⁾ was built in this way, using $G = SU(2)_L \otimes SU(2)_R$, $H = SU(2)_V$, and considering the gauging of $SU(2)_W \otimes U(1)_Y$.

The model with vector and axial vector resonances can be built using the same technique. In particular I will present the results for the case in which the new resonances are degenerate in mass (neglecting the weak corrections).

This type of realization corresponds to a maximal symmetry $[SU(2) \otimes SU(2)]^3$. After gauging the standard $SU(2)_W \otimes U(1)_Y$, the model describes the ordinary gauge bosons $W^\pm$, Z and γ and, in addition, two new triplets of massive gauge bosons, $L^\pm$, L_3 , $R^\pm$, R_3 , self-interacting with gauge coupling constant g'' . These heavy resonances, as a consequence of the chiral symmetry, are degenerate in mass M in the $g'' \rightarrow \infty$ limit.

This degenerate model has the interesting feature of allowing for a strong electroweak resonant sector at relatively low energies, while satisfying the severe constraints from existing LEP, SLC and CDF data (see Fig. 1).




Fig. 1. 90% C.L. contour on the plane $(M, g/g'')$ obtained by comparing the values of the ϵ parameters from the model with the experimental data from LEP. The allowed region is below the curve.

2. Degenerate BESS at future e^+e^- colliders

In this section I will discuss the sensitivity of the model at LEP2 and future e^+e^- linear colliders, for different options of total centre of mass energies and luminosities. Cross-sections and asymmetries for the channel $e^+e^- \rightarrow f^+f^-$ and $e^+e^- \rightarrow W^+W^-$ in the Standard Model and in the degenerate BESS model at tree level have been studied. In the fermion channel the study is based on the following observables: the total hadronic ($\mu^+\mu^-$) cross-sections σ^h (σ^μ), the forward-backward and left-right asymmetries $A_{FB}^{e^+e^- \rightarrow \mu^+\mu^-}$, $A_{FB}^{e^+e^- \rightarrow b\bar{b}}$, $A_{LR}^{e^+e^- \rightarrow \mu^+\mu^-}$, $A_{LR}^{e^+e^- \rightarrow h}$ and $A_{LR}^{e^+e^- \rightarrow b\bar{b}}$. At LEP2 we can add to

the previous observables the W mass measurement.

The result of this analysis shows that LEP2 will not improve considerably the existing limits ³⁾. To improve these limits it is necessary to consider higher energy colliders. Two options for a high energy e^+e^- collider have been studied: $\sqrt{s} = 500 \text{ GeV}$ ($\sqrt{s} = 1 \text{ TeV}$) with an integrated luminosity of $20fb^{-1}$ ($80fb^{-1}$). In Fig. 2 a combined picture of the 90% C.L. contours on the plane $(M, g/g'')$ from e^+e^- at two values of $\sqrt{s}$ is shown. The dotted line represents the limit from the combined unpolarized observables at $\sqrt{s} = 500 \text{ GeV}$ with an integrated luminosity of $20fb^{-1}$; the dashed line is the limit from the combined unpolarized observables at $\sqrt{s} = 1000 \text{ GeV}$ with an integrated luminosity of $80fb^{-1}$. As expected increasing the energy of the collider and rescaling the integrated luminosity gives stronger bounds on the parameter space.

In conclusion a substantial improvement with respect to the LEP bounds, even without polarized beams, is obtained. Considering the polarized observables one gets a further improvement on the limits. The WW final state does not modify the strong limits obtained using the fermion final state. This is because the degenerate model has no strong enhancement of the WW channel, present in the usual strong electroweak models.




Fig. 2. 90% C.L. contour on the plane $(M, g/g'')$ from e^+e^- at $\sqrt{s} = 500 \text{ GeV}$ with an integrated luminosity of $20fb^{-1}$ and $\sqrt{s} = 1000 \text{ GeV}$ with an integrated luminosity of $80fb^{-1}$. Allowed regions are below the curves.

References

- [1] R. Casalbuoni, A. Deandrea, S. De Curtis, D. Dominici, R. Gatto and M. Grazzini, UGVA-DPT 1995/10-906, hep-ph/9510431.
- [2] R. Casalbuoni, S. De Curtis, D. Dominici and R. Gatto, Phys. Lett. **B155** (1985) 95;

Nucl. Phys. **B282** (1987) 235.

- [3] R. Casalbuoni, A. Deandrea, S. De Curtis, D. Dominici, R. Gatto and M. Grazzini, Proceedings of the CERN Workshop on Physics at Lep2, CERN Yellow Report, to appear in 1996.